PAST EXAM PAPER & MEMO N6

ABOUT THE QUESTION PAPERS AND ONLINE INSTANT ACCESS:

THANK YOU FOR DOWNLOADING THE PAST EXAM PAPER AND ITS MEMO, WE HOPE IT WILL BE OF HELP TO YOU. SHOULD YOU NEED MORE QUESTION PAPERS AND THEIR MEMOS PLEASE SEND US AN EMAIL TO INFO@ PREVIOUSPAPERS.CO.ZA AND WE WILL SEND YOU A FULL DOWNLOAD VERSION OF THESE THOUGH A FULL DOWNLOAD COMES AT A COST.

ALTERNATIVELY YOU CAN PURCHASE THEM ONLINE FROM OUR WEBSITE WWW.PREVIOUSPAPERS.CO.ZA TO AVOID DELAYS AND YOU FOLLOW ALL THE NEEDED STEPS TO GET INSTANT DOWNLOAD. **WITH ONLINE PAYMENT YOU CAN GET INSTANT DOWNLOADS OF YOUR PURCHASE**.

WE SELL PREVIOUS PAPERS AND MEMOS FOR THE SUBJECTS MENTIONED AND THE PAPERS ARE BETWEEN 2014-2019. THE PAPERS ARE IN PDF FORM AND EACH PDF HAS A MINIMUM OF SEVEN DIFFERENT PAPERS. THE YEARS FOR THE PAPERS YOU ARE PURCHASING ARE ALSO INCLUDED ON THE WEBSITE.

PRICE OF THE PAPERS AT A BIG DISCOUNT

Previous papers are very important in ensuring you pass your final exams. The actual value of the papers we are making you access is way more than R1 000 and we are making these very cheap as we have you in mind. For a small amount you can have these papers.

THE COST PER SUBJECT IS R300.

higher education & training

Department:
Higher Education and Training
REPUBLIC OF SOUTH AFRICA

T980**(E)**(A6)T

APRIL EXAMINATION

NATIONAL CERTIFICATE

MATHEMATICS N6

(16030186)

6 April 2016 (X-Paper)
09:00–12:00

Calculators may be used.

This question paper consists of 5 pages and 1 formula sheet of 7 pages.

Copyright reserved

DEPARTMENT OF HIGHER EDUCATION AND TRAINING
REPUBLIC OF SOUTH AFRICA

NATIONAL CERTIFICATE
MATHEMATICS N6
TIME: 3 HOURS
MARKS: 100

INSTRUCTIONS AND INFORMATION

1. Answer ALL the questions.
2. Read ALL the questions carefully.
3. Number the answers according to the numbering system used in this question paper.
4. Questions may be answered in any order, but subsections of questions must be kept together.
5. Show ALL the intermediate steps.
6. ALL the formulae used must be written down.
7. Questions must be answered in BLUE or BLACK ink.
8. Write neatly and legibly.

Copyright reserved

QUESTION 1

1.1 If $z = -5x^3y^2 - y^4 + 3x^2y$, determine $\frac{\partial^2 z}{\partial x \partial y}$ (2)

1.2 Given: $I = \frac{V}{R}$

Calculate the change in I if V decreases with 5 volts and R with 8 ohms. The original value of V is 30 volts and of R is 10 ohms. (4)
[6]

QUESTION 2

Determine $\int y\,dx$ if:

2.1 $y = \sin^4 5x \cos^3 5x$ (5)

2.2 $y = \frac{1}{\sqrt{16x - x^2}}$ (3)

2.3 $y = \sin^4 mx$ (4)

2.4 $y = e^{\frac{x}{2}}.\cos 3x$ (6)
[18]

QUESTION 3

Use partial fractions to calculate the following integrals:

3.1 $\int \frac{-x^2 + 3x + 4}{x(1-2x)^2}\,dx$ (6)

3.2 $\int \frac{10x^2 + 7x + 1}{(2x^2+1)(4x-1)}\,dx$ (6)
[12]

Copyright reserved

QUESTION 4

4.1 Calculate the particular solution of:

$$2\sin x\frac{dy}{dx} - y(\sin 2x) = \frac{2\sin x}{\sec x} \quad \text{at} \quad (0;1)$$

(5)

4.2 Calculate the particular solution of:

$$\frac{d^2y}{dx^2} - 7\frac{dy}{dx} + 6y = 2x + 3, \text{ if } y = 1 \text{ when } x = 0 \text{ and } \frac{dy}{dx} = 2 \text{ when } x = 0.$$

(7)
[12]

QUESTION 5

5.1 5.1.1 Calculate the points of intersection of the two curves $y = \frac{3}{x}$ and $y + x - 4 = 0$. Make a neat sketch of the curves and show the area, in the first quadrant, bounded by the curves. Show the representative strip/element that you will use to calculate the volume (use the SHELL method only) generated if the area bounded by the curves rotates about the y-axis. (3)

5.1.2 Use the SHELL method to calculate the volume generated if the area, described in QUESTION 5.1.1, bounded by the two curves $y = \frac{3}{x}$ and $y + x - 4 = 0$, rotates about the y-axis. (5)

5.2 5.2.1 Make a neat sketch of the graph $y = \tan x$. Show the representative strip/element that you will use to calculate the volume generated if the area bounded by the graph, the ordinates $y = 0$ and $x = \frac{\pi}{3}$ rotates about the x-axis. (2)

5.2.2 Calculate the volume generated if the area, described in QUESTION 5.1.1, rotates about the x-axis. (3)

5.2.3 Calculate the volume moment about the y-axis as well as the distance of the centre of gravity from the y-axis. (6)

5.3 5.3.1 Calculate the points of intersection of the two curves $y = 2x^2$ and $x = \frac{y}{3}$. Make a neat sketch of the curves and show the area bounded by the curves. Show the representative strip/element, PERPENDICULAR to the x-axis, that you will use to calculate the area bounded by the curves. (3)

Copyright reserved

5.3.2 Calculate the area described in QUESTION 5.3.1, bounded by the two curves $y = 2x^2$ and $x = \frac{y}{3}$. (3)

5.3.3 Calculate the second moment of area of the area described in QUESTION 5.3.1 about the *y*-axis. (4)

5.3.4 Express the answer in QUESTION 5.3.3 in terms of the area. (1)

5.4 5.4.1 A weir in the form of a trapezium is 2 m high, 10 m wide at the top and 4 m wide at the bottom. The top of the weir is in the water surface.

Sketch the weir and show the representative strip/element that you will use to calculate the depth of the centre of pressure on the retaining wall.

Calculate the relation between the two variables *x* and *y*. (3)

5.4.2 Calculate, by using integration, the area moment of the weir about the water level. (3)

5.4.3 Calculate, by using integration, the second moment of area of the weir about the water level, as well as the depth of the centre of pressure on the weir. (4)

[40]

QUESTION 6

6.1 Calculate the arc length of the curve described by the parametric equations, $x = 5(\cos t + t\sin t)$ and $y = 5(\sin t - t\cos t)$, between the points $t = 0$ and $t = \pi$. (6)

6.2 Calculate the surface area generated when the curve of $y = \sqrt{16x}$, over the interval $1 \le x \le 4$, is rotated about the *x*-axis. (6)

[12]

TOTAL: 100

Copyright reserved

MATHEMATICS N6

FORMULA SHEET

Any other applicable formula may also be used.

Trigonometry

$\sin^2 x + \cos^2 x = 1$

$1 + \tan^2 x = \sec^2 x$

$1 + \cot^2 x = \operatorname{cosec}^2 x$

$\sin 2A = 2 \sin A \cos A$

$\cos 2A = \cos^2 A - \sin^2 A$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

$\sin^2 A = ½ - ½ \cos 2A$

$\cos^2 A = ½ + ½ \cos 2A$

$\sin (A \pm B) = \sin A \cos B \pm \sin B \cos A$

$\cos (A \pm B) = \cos A \cos B \mp \sin A \sin B$

$$\tan (A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$\sin A \cos B = ½ [\sin (A + B) + \sin (A - B)]$

$\cos A \sin B = ½ [\sin (A + B) - \sin (A - B)]$

$\cos A \cos B = ½ [\cos (A + B) + \cos (A - B)]$

$\sin A \sin B = ½ [\cos (A - B) - \cos (A + B)]$

$$\tan x = \frac{\sin x}{\cos x} ; \sin x = \frac{1}{\operatorname{cosec} x} ; \cos x = \frac{1}{\sec x}$$

Copyright reserved

$f(x)$	$\frac{d}{dx} f(x)$	$\int f(x)dx$
x^n	nx^{n-1}	$\frac{x^{n+1}}{n+1} + C \quad (n \neq -1)$
ax^n	$a \frac{d}{dx} x^n$	$a \int x^n \, dx$
e^{ax+b}	$e^{ax+b} . \frac{d}{dx} (ax + b)$	$\frac{e^{ax+b}}{\frac{d}{dx}(ax+b)} + C$
a^{dx+e}	$a^{dx+e} . \ln a . \frac{d}{dx} (dx + e)$	$\frac{a^{dx+e}}{\ln a . \frac{d}{dx}(dx+e)} + C$
$\ln(ax)$	$\frac{1}{ax} . \frac{d}{dx} ax$	$x \ln ax - x + C$
$e^{f(x)}$	$e^{f(x)} \frac{d}{dx} f(x)$	-
$a^{f(x)}$	$a^{f(x)} . \ln a . \frac{d}{dx} f(x)$	-
$\ln f(x)$	$\frac{1}{f(x)} . \frac{d}{dx} f(x)$	-
$\sin ax$	$a \cos ax$	$-\frac{\cos ax}{a} + C$
$\cos ax$	$-a \sin ax$	$\frac{\sin ax}{a} + C$
$\tan ax$	$a \sec^2 ax$	$\frac{1}{a} \ln [\sec (ax)] + C$
$\cot ax$	$-a \operatorname{cosec}^2 ax$	$\frac{1}{a} \ln [\sin (ax)] + C$
$\sec ax$	$a \sec ax \tan ax$	$\frac{1}{a} \ln [\sec ax + \tan ax] + C$
$\operatorname{cosec} ax$	$-a \operatorname{cosec} ax \cot ax$	$\frac{1}{a} \ln \left[\tan \left(\frac{ax}{2} \right) \right] + C$

Copyright reserved

$f(x)$	$\frac{d}{dx} f(x)$	$\int f(x)\,dx$
$\sin f(x)$	$\cos f(x) \,.\, f'(x)$	-
$\cos f(x)$	$-\sin f(x) \,.\, f'(x)$	-
$\tan f(x)$	$\sec^2 f(x) \,.\, f'(x)$	-
$\cot f(x)$	$-\text{cosec}^2 f(x) \,.\, f'(x)$	-
$\sec f(x)$	$\sec f(x) \tan f(x) \,.\, f'(x)$	-
$\text{cosec}\, f(x)$	$-\text{cosec}\, f(x) \cot f(x) \,.\, f'(x)$	-
$\sin^{-1} f(x)$	$\frac{f'(x)}{\sqrt{1-[f(x)]^2}}$	-
$\cos^{-1} f(x)$	$\frac{-f'(x)}{\sqrt{1-[f(x)]^2}}$	-
$\tan^{-1} f(x)$	$\frac{f'(x)}{[f(x)]^2+1}$	-
$\cot^{-1} f(x)$	$\frac{-f'(x)}{[f(x)]^2+1}$	-
$\sec^{-1} f(x)$	$\frac{f'(x)}{f(x)\sqrt{[f(x)]^2-1}}$	-
$\text{cosec}^{-1} f(x)$	$\frac{-f'(x)}{f(x)\sqrt{[f(x)]^2-1}}$	-
$\sin^2(ax)$	-	$\frac{x}{2} - \frac{\sin(2ax)}{4a} + C$
$\cos^2(ax)$	-	$\frac{x}{2} + \frac{\sin(2ax)}{4a} + C$
$\tan^2(ax)$	-	$\frac{1}{a}\tan(ax) - x + C$

Copyright reserved

$f(x)$	$\frac{d}{dx} f(x)$	$\int f(x)\,dx$
$\cot^2(ax)$	-	$-\frac{1}{a}\cot(ax) - x + C$

$$\int f(x)\,g'(x)\,dx = f(x)\,g(x) - \int f'(x)\,g(x)\,dx$$

$$\int [f(x)]^n\, f'(x)\,dx = \frac{[f(x)]^{n+1}}{n+1} + C \qquad (n \neq -1)$$

$$\int \frac{f'(x)}{f(x)}\,dx = \ln f(x) + C$$

$$\int \frac{dx}{\sqrt{a^2 - b^2x^2}} = \frac{1}{b}\sin^{-1}\frac{bx}{a} + C$$

$$\int \frac{dx}{a^2 + b^2x^2} = \frac{1}{ab}\tan^{-1}\frac{bx}{a} + C$$

$$\int \sqrt{a^2 - b^2x^2}\,dx = \frac{a^2}{2b}\sin^{-1}\frac{bx}{a} + \frac{x}{2}\sqrt{a^2 - b^2x^2} + C$$

$$\int \frac{dx}{a^2 - b^2x^2} = \frac{1}{2ab}\ln\left(\frac{a + bx}{a - bx}\right) + C$$

$$\int \sqrt{x^2 \pm b^2}\,dx = \frac{x}{2}\sqrt{x^2 \pm b^2} \pm \frac{b^2}{2}\ln\left[x + \sqrt{x^2 \pm b^2}\right] + C$$

$$\int \frac{dx}{\sqrt{b^2x^2 \pm a^2}} = \frac{1}{b}\ln\left[bx + \sqrt{b^2x^2 \pm a^2}\right] + C$$

Applications of integration

AREAS

$$A_x = \int_a^b y\,dx\,;\; A_x = \int_a^b (y_1 - y_2)\,dx$$

$$A_y = \int_a^b x\,dy\,;\; A_y = \int_a^b (x_1 - x_2)\,dy$$

Copyright reserved

VOLUMES

$$V_x = \pi\int_a^b y^2dx\,;\, V_x = \pi\int_a^b \left(y_1^2 - y_2^2\right)dx\,;\, V_x = 2\pi\int_a^b xydy$$

$$V_y = \pi\int_a^b x^2dy\,;\, V_y = \pi\int_a^b \left(x_1^2 - x_2^2\right)dy\,;\, V_y = 2\pi\int_a^b xydx$$

AREA MOMENTS

$$A_{m-x} = rdA \qquad A_{m-y} = rdA$$

CENTROID

$$\bar{x} = \frac{A_{m-y}}{A} = \frac{\int_a^b rdA}{A}\,;\, \bar{y} = \frac{A_{m-x}}{A} = \frac{\int_a^b rdA}{A}$$

SECOND MOMENT OF AREA

$$I_x = \int_a^b r^2dA \quad ; \quad I_y = \int_a^b r^2dA$$

VOLUME MOMENTS

$$V_{m-x} = \int_a^b rdV \quad ; \quad V_{m-y} = \int_a^b rdV$$

CENTRE OF GRAVITY

$$\bar{x} = \frac{v_{m-y}}{V} = \frac{\int_a^b rdV}{V}\;;\quad \bar{y} = \frac{v_{m-x}}{V} = \frac{\int_a^b rdV}{V}$$

MOMENTS OF INERTIA

Mass = Density × volume

$M = \rho V$

DEFINITION: $I = m\,r^2$

Copyright reserved

GENERAL

$$I = \int_a^b r^2 dm = \rho \int_a^b r^2 dV$$

CIRCULAR LAMINA

$$I_z = \frac{1}{2} mr^2$$

$$I = \frac{1}{2} \int_a^b r^2 dm = \frac{1}{2} \rho \int_a^b r^2 dV$$

$$I_x = \frac{1}{2} \rho\pi \int_a^b y^4 dx \qquad I_y = \frac{1}{2} \rho\pi \int_a^b x^4 dy$$

CENTRE OF FLUID PRESSURE

$$\bar{\bar{y}} = \frac{\int_a^b r^2 dA}{\int_a^b r dA}$$

$$\frac{f(x)}{(ax+b)^n} = \frac{A}{ax+b} + \frac{B}{(ax+b)^2} + \frac{C}{(ax+b)^3} + \ldots \frac{Z}{(ax+b)^n}$$

$$\frac{f(x)}{(ax+b)^3 (cx+d)^3} = \frac{A}{ax+b} + \frac{B}{(ax+b)^2} + \frac{C}{(ax+b)^3} + \frac{D}{(cx+d)} + \frac{E}{(cx+d)^2} + \frac{F}{(cx+d)^3}$$

$$\frac{f(x)}{(ax^2+bx+c)(dx+e)^n} = \frac{Ax+F}{ax^2+bx+c} + \frac{B}{dx+e} + \frac{C}{(dx+e)^2} + \ldots + \frac{Z}{(dx+e)^n}$$

$$A_x = \int_a^b 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2}\, dx$$

$$A_x = \int_d^c 2\pi y \sqrt{1 + \left(\frac{dx}{dy}\right)^2}\, dy$$

$$A_y = \int_a^b 2\pi x \sqrt{1 + \left(\frac{dy}{dx}\right)^2}\, dx$$

$$A_y = \int_d^c 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2}\, dy$$

Copyright reserved

$$A_x = \int_{u1}^{u2} 2\pi y \sqrt{\left(\frac{dx}{du}\right)^2 + \left(\frac{dy}{du}\right)^2}\, du$$

$$A_y = \int_{u1}^{u2} 2\pi x \sqrt{\left(\frac{dx}{du}\right)^2 + \left(\frac{dy}{du}\right)^2}\, du$$

$$S = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2}\, dx$$

$$S = \int_c^d \sqrt{1 + \left(\frac{dx}{dy}\right)^2}\, dy$$

$$S = \int_{u1}^{u2} \sqrt{\left(\frac{dx}{du}\right)^2 + \left(\frac{dy}{du}\right)^2}\, du$$

$$\frac{dy}{dx} + Py = Q \quad \therefore\ ye^{\int Pdx} = \int Qe^{\int Pdx}\, dx$$

$$y = Ae^{r_1 x} + Be^{r_2 x} \quad r_1 \neq r_2$$

$$y = e^{rx}(A + Bx) \quad r_1 = r_2$$

$$y = e^{ax}[A\cos bx + B\sin bx] \quad r = a \pm ib$$

$$\frac{d^2y}{dx^2} = \frac{d}{d\theta}\left(\frac{dy}{dx}\right)\frac{d\theta}{dx}$$

Copyright reserved

higher education & training

Department:
Higher Education and Training
REPUBLIC OF SOUTH AFRICA

MARKING GUIDELINE

NATIONAL CERTIFICATE

APRIL EXAMINATION

MATHEMATICS N6

6 APRIL 2016

This marking guideline consists of 17 pages.

Copyright reserved

TOTAL: $\frac{200}{2} = 100$

NOTE: Do NOT subtract marks for incorrect units or units omitted.

QUESTION 1

1.1 $z = -5x^3y^2 - y^4 + 3x^2y$

$$\frac{\partial}{\partial x}\left(\frac{\partial z}{\partial y}\right) = \frac{\partial}{\partial x}(-10x^3y - 4y^3 + 3x^2) \quad \checkmark \quad \checkmark \quad \checkmark$$

$$= -30x^2y + 6x \quad \checkmark$$

(4)

1.2 $I = \frac{V}{R}$

$I = VR^{-1}$

$$\Delta I = \frac{\partial I}{\partial V}\Delta V + \frac{\partial I}{\partial R}\Delta R \quad \checkmark \quad \checkmark$$

$$= R^{-1}\Delta V - VR^{-2}\Delta R \quad \checkmark \quad \checkmark$$

$$= \frac{1}{R}\Delta V - \frac{V}{R^2}\Delta R \quad \checkmark$$

$$= \frac{1}{(10)}(-5) - \frac{30}{(10)^2}(-8) \quad \checkmark \quad \checkmark$$

$= 1{,}9$ A $\checkmark$

(8)
[12]

QUESTION 2

2.1 $y = \int \sin^4 5x \cos^3 5x dx$

✓

$= \int \sin^4 5x \cos^2 5x . \cos 5x dx$

✓

$= \int \sin^4 5x(1-\sin^2 5x)\cos 5x dx$

$= \frac{1}{5}\int \sin^4 5x(1-\sin^2 5x)5\cos 5x dx$ $u = \sin 5x$ $du = 5\cos 5x dx$

✓ ✓ ✓

$= \frac{1}{5}\int u^4(1-u^2)du$

✓

$= \frac{1}{5}\int u^4 - u^6)du$

✓ ✓

$= \frac{1}{5}\left[\frac{u^5}{5} - \frac{u^7}{7}\right] + c$

✓ ✓

$= \frac{1}{5}\left[\frac{\sin^5 5x}{5} - \frac{\sin^7 5x}{7}\right] + c$

OR

$= \frac{1}{25}\sin^5 5x - \frac{1}{35}\sin^7 5x + c$

OR

$y = \int \sin^4 5x \cos^3 5x dx$

✓

$= \int \sin^4 5x \cos^2 5x . \cos 5x dx$

✓

$= \int \sin^4 5x(1-\sin^2 5x)\cos 5x dx$

✓ ✓ ✓ ✓

$= \int \sin^4 5x . \cos 5x dx - \int \sin^6 5x . \cos 5x dx$ $u = \sin 5x$ $du = 5\cos 5x dx$

✓ ✓ ✓ ✓

$= \frac{1}{5}.\frac{\sin^5 5x}{5} - \frac{1}{5}.\frac{\sin^7 5x}{7} + c$ (10)

2.2 $y = \int \frac{1}{\sqrt{16x - x^2}} dx$

$= -x^2 + 16x$

$= -(x^2 - 16x)$ ✓

✓ ✓

$= -[(x-8)^2 - 64]$

$= 64 - (x-8)^2$ ✓

$= \int \frac{1}{\sqrt{64 - (x-8)^2}} dx$ ✓

$= \sin^{-1} \frac{(x-8)}{8} + c$ ✓ (6)

2.3 $y = \int \sin^4 mx dx$

$= \int (\sin^2 mx)^2 dx$ ✓

$= \int (\frac{1}{2} - \frac{1}{2}\cos 2mx)^2 dx$ ✓

$= \int (\frac{1}{2} - \frac{1}{2}\cos 2mx)(\frac{1}{2} - \frac{1}{2}\cos 2mx) dx$

$= \int (\frac{1}{4} - \frac{1}{2}\cos 2mx + \frac{1}{4}\cos^2 2mx) dx$ ✓

✓ ✓ ✓ ✓ ✓

$= \frac{1}{4}x - \frac{1}{2} \cdot \frac{\sin 2mx}{2m} + \frac{1}{4}\left[\frac{x}{2} + \frac{\sin 4mx}{8m}\right] + c$

$= \frac{1}{4}x - \frac{\sin 2mx}{4m} + \frac{x}{8} + \frac{\sin 4mx}{32m} + c$ (8)

2.4 $y = \int e^{\frac{x}{2}}.\cos 3x\,dx$

✓ ✓

$\int y\,dx = e^{\frac{x}{2}}(\frac{\sin 3x}{3}) - \int \frac{1}{2}e^{\frac{x}{2}}(\frac{\sin 3x}{3})dx$

$f(x) = e^{\frac{x}{2}}$ $g'(x) = \cos 3x$

$f'(x) = \frac{1}{2}e^{\frac{x}{2}}$ $g(x) = \frac{\sin 3x}{3}$

✓

$= \frac{1}{3}e^{\frac{x}{2}}\sin 3x - \frac{1}{6}\int e^{\frac{x}{2}}.\sin 3x\,dx$

$f(x) = e^{\frac{x}{2}}$ $g'(x) = \sin 3x$

$f'(x) = \frac{1}{2}e^{\frac{x}{2}}$ $g(x) = -\frac{\cos 3x}{3}$

✓ ✓

$= \frac{1}{3}e^{\frac{x}{2}}\sin 3x - \frac{1}{6}\left[e^{\frac{x}{2}}.-\frac{\cos 3x}{3} - \int \frac{1}{2}e^{\frac{x}{2}}.-\frac{\cos 3x}{3}dx\right]$

✓ ✓

$= \frac{1}{3}e^{\frac{x}{2}}\sin 3x + \frac{1}{18}e^{\frac{x}{2}}.\cos 3x - \frac{1}{36}\int e^{\frac{x}{2}}.\cos 3x\,dx$

$I = \frac{1}{3}e^{\frac{x}{2}}\sin 3x + \frac{1}{18}e^{\frac{x}{2}}.\cos 3x - \frac{1}{36}I$ ✓

✓

$\therefore \frac{37}{36}I = \frac{1}{3}e^{\frac{x}{2}}\sin 3x + \frac{1}{18}e^{\frac{x}{2}}.\cos 3x$ ✓

✓ ✓

$I = \frac{36}{37}(\frac{1}{3}e^{\frac{x}{2}}\sin 3x + \frac{1}{18}e^{\frac{x}{2}}.\cos 3x) + c$

$= 0{,}973(0{,}333e^{\frac{x}{2}}\sin 3x + 0{,}054e^{\frac{x}{2}}.\cos 3x) + c$

$= 0{,}324e^{\frac{x}{2}}\sin 3x + 0{,}054e^{\frac{x}{2}}.\cos 3x + c$

OR

$y = \int e^{\frac{x}{2}}.\cos 3x\,dx$

✓ ✓

$\int y\,dx = 2e^{\frac{x}{2}}\cos 3x - \int(-3\sin 3x)2e^{\frac{x}{2}}dx$

$f(x) = \cos 3x$ $g'(x) = e^{\frac{x}{2}}$

$f'(x) = -3\sin 3x$ $g(x) = \frac{e^{\frac{x}{2}}}{\frac{1}{2}} = 2e^{\frac{x}{2}}$

✓

$= 2e^{\frac{x}{2}}\cos 3x + 6\int \sin 3x.e^{\frac{x}{2}}dx$

✓ ✓

$= 2e^{\frac{x}{2}}\cos 3x + 6\left[\sin 3x.2e^{\frac{x}{2}} - \int 3\cos 3x.2e^{\frac{x}{2}}dx\right]$

$f(x) = \sin 3x$ $g'(x) = e^{\frac{x}{2}}$

$f'(x) = 3\cos 3x$ $g(x) = 2e^{\frac{x}{2}}$

✓ ✓

$= 2e^{\frac{x}{2}}\cos 3x + 12\sin 3x.e^{\frac{x}{2}} - 36\int \cos 3x.e^{\frac{x}{2}}dx$

$$I = 2e^{\frac{x}{2}}\cos 3x + 12e^{\frac{x}{2}}.\sin 3x - 36I \quad \checkmark$$
$\checkmark$

$$\therefore 37.I = 2e^{\frac{x}{2}}\cos 3x + 12e^{\frac{x}{2}}.\sin 3x \quad \checkmark$$
$\checkmark \qquad \checkmark$

$$I = \frac{1}{37}(2e^{\frac{x}{2}}\cos 3x + 12e^{\frac{x}{2}}.\sin 3x) + c$$

$$= 0{,}054e^{\frac{x}{2}}\cos 3x + 0{,}324e^{\frac{x}{2}}\sin 3x + c \qquad (12)$$

[36]

QUESTION 3

3.1 $\int \frac{-x^2+3x+4}{x(1-2x)^2}\,dx$

$$\frac{-x^2+3x+4}{x(1-2x)^2} = \frac{A}{x} + \frac{B}{(1-2x)^2} + \frac{C}{(1-2x)} \quad \checkmark$$

$$-x^2+3x+4 = A(1-2x)^2 + Bx + Cx(1-2x) \quad \checkmark$$

$$Let \;\; x = 0; \quad \therefore A = 4 \quad \checkmark$$

$$Let \;\; x = \frac{1}{2}; \quad \therefore B = \frac{21}{2} \quad (10{,}5) \quad \checkmark$$

$$-x^2+3x+4 = A - 4Ax + 4Ax^2 + Bx + Cx - 2Cx^2 \quad \checkmark$$

Equate coeff of x^2: $\quad C = \frac{17}{2} \quad (8{,}5) \quad \checkmark$

$$= \int \frac{4}{x}dx + \int \frac{\frac{21}{2}}{(1-2x)^2}dx + \int \frac{\frac{17}{2}}{(1-2x)}dx \quad \checkmark$$
$\checkmark \quad \checkmark \quad \checkmark \quad \checkmark \quad \checkmark$

$$= 4\ln x - \frac{21}{4}.\frac{(1-2x)^{-1}}{-1} - \frac{17}{4}\ln(1-2x) + c$$

$$= 4\ln x + \frac{21}{4(1-2x)} - \frac{17}{4}\ln(1-2x) + c$$

$$= 0{,}444\ln x + \frac{5{,}25}{(1-2x)} - 1{,}444\ln(1-2x) + c \qquad (12)$$

3.2 $\int \frac{10x^2+7x+1}{(2x^2+1)(4x-1)}\,dx$

$$\frac{10x^2+7x+1}{(2x^2+1)(4x-1)} = \frac{Ax+B}{2x^2+1} + \frac{C}{4x-1} \quad \checkmark$$

$$10x^2+7x+1 = (Ax+B)(4x-1) + C(2x^2+1) \quad \checkmark$$

$$10x^2+7x+1 = 4Ax^2+4Bx-Ax-B+2Cx^2+C \quad \checkmark$$

$$let \quad x = \frac{1}{4} \qquad \therefore C = 3 \quad \checkmark$$

Equate coeff of x^2: $A = 1$ ✓ *Equate* x: $B = 2$ ✓

$$= \int \frac{x+2}{2x^2+1}dx + \int \frac{3}{4x-1}dx \quad \checkmark$$

$$= \int \frac{x}{2x^2+1}dx + \int \frac{2}{2x^2+1}dx + \int \frac{3}{4x-1}dx \quad \checkmark$$

$$= \frac{1}{4}\ln(2x^2+1) + 2\left(\frac{1}{\sqrt{2}}\right)\arctan\sqrt{2}x + \frac{3}{4}\ln(4x-1) + c \quad \checkmark$$

(12)
[24]

QUESTION 4

4.1 $2\sin x\frac{dy}{dx} - y(\sin 2x) = \frac{2\sin x}{\sec x}$

$$\frac{dy}{dx} - \frac{y(2\sin 2x)}{2\sin x} = \frac{1}{\sec x} \quad \checkmark$$

$$\frac{dy}{dx} - \frac{y(2\sin x\cos x)}{2\sin x} = \frac{1}{\sec x} \quad \checkmark$$

$$\frac{dy}{dx} - y.\cos x = \cos x \quad \checkmark$$

$$e^{\int -\cos x.dx} = e^{-\sin x} \quad \checkmark$$

$$e^{-\sin x}.y = \int e^{-\sin x}.\cos x dx \quad \checkmark$$

$$= -e^{-\sin x} + c \quad \checkmark$$

$$e^{-\sin 0}.(1) = -e^{-\sin 0} + c \quad \checkmark$$

$$\therefore c = 2 \quad \checkmark$$

$$e^{-\sin x}.y = -e^{-\sin x} + 2 \quad \checkmark$$

(10)

4.2 $\frac{d^2y}{dx^2} - 7\frac{dy}{dx} + 6y = 2x + 3$

y_c : $m^2 - 7m + 6 = 0$ ✓

$(m-6)(m-1) = 0$

$m = 6; \quad m = 1$

$y_c = Ae^{6x}$ ✓

To find y_p $\therefore y = Cx + D$ ✓

$\frac{dy}{dx} = C$ ✓

$\frac{d^2y}{dx^2} = 0$ ✓

$0 - 7C + 6Cx + 6D = 2x + 3$ ✓

$C = \frac{1}{3}$ (0,333) ✓

$-7C + 6D = 3 \quad \therefore D = \frac{8}{9}$ (0,889) ✓

$\therefore y_p = \frac{1}{3}x + \frac{8}{9}$

$y = Ae^{6x} + Be^{x} + \frac{1}{3}x + \frac{8}{9}$ ✓

$1 = A + B + \frac{8}{9} \quad \therefore A + B = \frac{1}{9}$ ✓

$\frac{dy}{dx} = 6Ae^{6x} + Be^{x} + \frac{1}{3}$ ✓

$2 = 6A + B + \frac{1}{3}$

$\therefore A = 0,311 \quad \left(\frac{14}{45}\right)$ ✓

$and \quad B = -0,2 \quad \left(-\frac{9}{45} or -\frac{1}{5}\right)$ ✓

$\therefore y = \frac{14}{45}e^{6x} - \frac{1}{5}e^{x} + \frac{1}{3}x + \frac{8}{9}$ ✓

$y = 0,311e^{6x} - 0,2e^{x} + 0,333x + 0,889$

(14)
[24]

QUESTION 5

5.1 5.1.1

$$\frac{3}{x} = -x + 4$$

$$x^2 - 4x + 3 = 0$$

$$(x-3)(x-1) = 0$$

$$x = 3; \quad x = 1 \quad ✓$$

$$y = 1; \quad y = 3 \quad ✓ \quad \therefore (3;1) \text{ and } (1;3)$$



(6)

✓ ✓ ✓

5.1.2 $\Delta V_y = 2\pi x \times (y_2 - y_1) \times \Delta x$

✓

$$V_y = 2\pi \int_1^3 x(y_2 - y_1)dx$$

Incorrect limits: max 7 marks

✓ ✓

$$= 2\pi \int_1^3 x(-x + 4 - \frac{3}{x})dx$$

$$= 2\pi \int_1^3 (-x^2 + 4x - 3)dx \quad ✓$$

$$= 2\pi \left[-\frac{x^3}{3} + \frac{4x^2}{2} - 3x \right]_1^3 \quad ✓$$

$$= 2\pi \left\{ \left[-\frac{(3)^3}{3} + 2(3)^2 - 3(3) \right] - \left[-\frac{(1)^3}{3} + 2(1)^2 - 3(1) \right] \right\} \quad ✓$$

$$= 2{,}667\pi \ \ units^3 \ \ or \ \ 8{,}278 \ \ or \ \ \frac{8}{3}\pi \ \ units^3 \quad ✓$$

(10)

5.2 5.2.1



(4)

5.2.2 $\Delta V_x = \pi y^2 \Delta x$ ✓

✓

$V_x = \pi \int_0^{\frac{\pi}{3}} y^2 \, dx$

$= \pi \int_0^{\frac{\pi}{3}} (\tan^2 x) \, dx$ ✓

$= \pi \left[(\tan x - x) \right]_0^{\frac{\pi}{3}}$ ✓

$= \pi \left[(\tan \frac{\pi}{3} - \frac{\pi}{3}) - (\tan 0 - 0) \right]$ ✓

$= 0{,}685\pi \quad units^3 \quad or \quad 2{,}152 \, units^3$ ✓

Incorrect limits: max 3 marks

(6)

5.2.3 $\Delta M_y = \pi y^2 \Delta x \times x$ ✓ ✓

Incorrect limits: max 8 marks

$\therefore M_y = \pi \int_0^{\frac{\pi}{3}} xy^2 dx$ ✓

$= \pi \int_0^{\frac{\pi}{3}} x(\tan^2 x)dx$ ✓

$f(x) = x$	$g'(x) = \tan^2 x$
$f'(x) = 1$	$g(x) = \tan x - x$

$= \pi \left\{ \left[x(\tan x - x) \right]_0^{\frac{\pi}{3}} - \int_0^{\frac{\pi}{3}} (\tan x - x)dx \right\}$ ✓ ✓

$= \pi \left\{ \left[x(\tan x - x) \right]_0^{\frac{\pi}{3}} - \left[\ln\sec x - \frac{x^2}{2} \right]_0^{\frac{\pi}{3}} \right\}$ ✓ ✓

$= \pi \left\{ \left[\frac{\pi}{3}(\tan\frac{\pi}{3} - \frac{\pi}{3}) \right] - \left[\ln\sec\frac{\pi}{3} - \frac{(\frac{\pi}{3})^2}{2} \right] \right\}$ ✓

$= 0{,}572\pi \quad units^3 \quad or \quad 1{,}798 \quad u^3$ ✓

$\bar{x} = \frac{1{,}798}{2{,}152}$ ✓

$= 0{,}836\, units$ ✓ (12)

5.3 5.3.1 $2x^2 = 3x$

$2x^2 - 3x = 0$

$x(2x - 3) = 0$

$x = 0; \quad x = \frac{3}{2}$ ✓

$y = 0; \quad y = 4\frac{1}{2}$ ✓ $\therefore (0;0)$ and $(\frac{3}{2};\frac{9}{2})$ (6)

5.3.2 $\Delta A = (y_2 - y_1)\Delta x$ ✓

✓

$A = \int_{\frac{1}{2}}^{1}(y_2 - y_1)dx$

Incorrect limits: max 3 marks

$= \int_{\frac{1}{2}}^{1}(3x - 2x^2)dx$ ✓

$= \left[\frac{3x^2}{2} - \frac{2x^3}{3}\right]_{\frac{1}{2}}^{1}$ ✓

$= \left[\frac{3(1)^2}{2} - \frac{2(1)^3}{3}\right] - \left[\frac{3(\frac{1}{2})^2}{2} - \frac{2(\frac{1}{2})^3}{3}\right]$ ✓

$= 0{,}542 \ units^2$ ✓ (6)

5.3.3 ✓ ✓

$\therefore \Delta I_y = (y_2 - y_1)\Delta x \times x^2$

✓ ✓

$= \int_{\frac{1}{2}}^{1}(3x - 2x^2)x^2 dx$

Incorrect limits: max 5 marks

$= \int_{\frac{1}{2}}^{1}(3x^3 - 2x^4)dx$ ✓

$= \left[\frac{3x^4}{4} - \frac{2x^5}{5}\right]_{\frac{1}{2}}^{1}$ ✓

$= \left[\frac{3(1)^4}{4} - \frac{2(1)^5}{5}\right] - \left[\frac{3(\frac{1}{2})^4}{4} - \frac{2(\frac{1}{2})^5}{5}\right]$ ✓

$= 0{,}316 \quad units^4$ ✓ (8)

5.3.4 $I = \frac{0{,}316}{0{,}542}$ ✓

$= 0{,}553 \quad A$ ✓ (2)

5.4 5.4.1



$$\frac{y-y_1}{x-x_1}=\frac{y_2-y_1}{x_2-x_1}$$

$$\frac{y-0}{x-2}=\frac{2-0}{5-2} \quad ✓$$

$$y=\frac{2}{3}(x-2)$$

$$\therefore x=\frac{3}{2}y+2 \quad ✓ \qquad \therefore dA=2(\frac{3}{2}y+2)dy \quad \text{or} \quad \therefore dA=(3y+4)dy$$

(6)

5.4.2

$$\int_0^2 r\,dA$$

✓ ✓ ✓

Incorrect limits: max 4 marks

$$=\int_0^2(2-y)(3y+4)dy$$

$$=\int_0^2(6y+8-3y^2-4y)dy \quad ✓$$

$$=\left[\frac{6y^2}{2}+8y-\frac{3y^3}{3}-\frac{4y^2}{2}\right]_0^2 \quad ✓$$

$$=\left[\frac{6(2)^2}{2}+8(2)-\frac{3(2)^3}{3}-\frac{4(2)^2}{2}\right]$$

$$=12 \text{ units}^3 \quad ✓$$

(6)

5.4.3 $\int_0^2 r^2 dA$

✓ ✓

Incorrect limits: max 5 marks

$= \int_0^2 (2-y)^2(3y+4)dy$

$= \int_0^2 (12y - 12y^2 + 3y^3 + 16 - 16y + 4y^2)dy$ ✓

$= \left[\frac{12y^2}{2} - \frac{12y^3}{3} + \frac{3y^4}{4} + 16y - \frac{16y^2}{2} + \frac{4y^3}{3}\right]_0^2$ ✓

$= \left[6(2)^2 - 4(2)^3 + \frac{3(2)^4}{4} + 16(2) - 8(2)^2 + \frac{4}{3}(2)^3\right]$ ✓

$= 14{,}667 \quad \text{units}^4$ ✓

$= y = \frac{14{,}667}{12}$ ✓

$= 1{,}222\, units$ ✓

(8)

Or alternative method

5.4 5.4.1



✓ ✓ ✓ ✓

$\frac{y-y_1}{x-x_1} = \frac{y_2-y_1}{x_2-x_1}$

$\frac{y-0}{x-5} = \frac{-2-0}{2-5}$

$y = \frac{2}{3}(x-5)$ ✓

$\therefore x = \frac{3}{2}y + 5$ ✓ $\quad \therefore dA = 2(\frac{3}{2}y + 5)dy$ or $\therefore dA = (3y + 10)dy$

(6)

5.4.2 ✓ ✓ ✓

$$= \int_{-2}^{0} y(3y+10)dy$$

Incorrect limits: max 4 marks

$$= \int_{-2}^{0} (3y^2+10y)dy \quad ✓$$

$$= \left[\frac{3y^3}{3}+\frac{10y^2}{2}\right]_{-2}^{0} \quad ✓$$

$$= \left[\frac{3(0)^3}{3}+\frac{10(0)^2}{2}\right]-\left[\frac{3(-2)^3}{3}+\frac{10(-2)^2}{2}\right] ✓$$

$$= -12 \text{ units}^3 \quad ✓$$ (6)

5.4.3 ✓ ✓

$$= \int_{-2}^{0} y^2(3y+10)dy$$

Incorrect limits: max 5 marks

$$= \int_{-2}^{0} (3y^3+10y^2)dy \quad ✓$$

$$= \left[\frac{3y^4}{4}+\frac{10y^3}{3}\right]_{-2}^{0} \quad ✓$$

$$= \left[\frac{3(0)^4}{4}+\frac{10(0)^3}{3}\right]-\left[\frac{3(-2)^4}{4}+\frac{10(-2)^3}{3}\right] ✓$$

$$= 14{,}667 \text{ units}^4 \quad ✓$$

$$\overline{y} = \frac{14{,}667}{-12} \quad ✓$$

$$= -1{,}222 \, units \quad ✓$$

(8)

[80]

QUESTION 6

6.1 $x = 5(\cos t + t\sin t)$ and $y = 5(\sin t - t\cos t)$

$\frac{dx}{dt} = 5(-\sin t + t\cos t + \sin t)$ ✓ $\frac{dy}{dt} = 5(\cos t + t\sin t - \cos t)$ ✓

Or $\frac{dx}{dt} = -5\sin t + 5t\cos t + 5\sin t$ OR $\frac{dy}{dt} = 5\cos t + 5t\sin t - 5\cos t$

$= 5t\cos t$ ✓ $= 5t\sin t$ ✓

$\left(\frac{dx}{dt}\right)^2 = (5t\cos t)^2$ $\left(\frac{dy}{dt}\right)^2 = (5t\sin t)^2$

$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = 25t^2\cos^2 t + 25t^2\sin^2 t$ ✓

$= 25t^2(\cos^2 t + \sin^2 t)$

$= 25t^2$ ✓

✓

$S = \int_0^{\pi} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\, dt$

Incorrect limits: max 9 marks

$= \int_0^{\pi} \sqrt{25t^2}\, dt$ ✓

$= 5\int_0^{\pi} t.dt$ ✓

$= 5\left[\frac{t^2}{2}\right]_0^{\pi}$ ✓

$= \frac{5}{2}\left[\pi^2\right]$ ✓

$= 2{,}5\pi^2$ units or 24,674 units ✓ (12)

6.2 $y = \sqrt{16x}$

$y = 4x^{\frac{1}{2}}$ ✓

$\frac{dy}{dx} = 2x^{-\frac{1}{2}}$ ✓

$\left(\frac{dy}{dx}\right)^2 = 4x^{-1}$ ✓

$1 + \left(\frac{dy}{dx}\right)^2 = 1 + \frac{4}{x}$ ✓

$= \frac{x+4}{x}$ ✓

✓

$A_x = 2\pi\int_1^4 y\sqrt{\frac{x+4}{x}}\, dx$ ✓

Incorrect limits: max 9 marks

✓

$$= 2\pi\int_1^4 4x^{\frac{1}{2}}\sqrt{\frac{x+4}{x}}dx$$

$$= 8\pi\int_1^4 \sqrt{x+4}dx \checkmark$$

$$= 8\pi\int_1^4 (x+4)^{\frac{1}{2}}dx$$

$$= 8\pi\left[\frac{(x+4)^{\frac{3}{2}}}{\frac{3}{2}}\right]_1^4 \checkmark$$

$$= \frac{16\pi}{3}\left[(x+4)^{\frac{3}{2}}\right]_1^4$$

$$= \frac{16\pi}{3}\left[(8)^{\frac{3}{2}} - (5)^{\frac{3}{2}}\right]_1^4 \checkmark$$

$= 61{,}051\ \pi$ units^2 or 191,798 units^2 ✓ (12)

[24]

TOTAL: 100

Copyright reserved

Website: **www.previouspapers.co.za**
Email: info@ previouspapers.co.za
CELL: 073 770 3028

THANK YOU FOR DOWNLOADING PAST EXAM PAPER & MEMO FOR N6

IF YOU NEED MORE PAPERS WE SELL THEM FOR R300

VISIT OUR WEBSITE AT WWW.PREVIOUSPAPERS.CO.ZA

OR SEND US AN EMAIL AT INFO@PREVIOUSPAPERS.CO.ZA AND

WE WILL GLADLY HELP YOU.